

Generic weak isolated horizons

Ayan Chatterjee* and Amit Ghosh†

Theory Division, Saha Institute of Nuclear Physics, Kolkata 700064, India

Weak isolated horizon boundary conditions have been relaxed supposedly to their weakest form such that both zeroth and the first law of black hole mechanics still emerge, thus making the formulation more amenable for applications in both analytic and numerical Relativity. As an additional gain it explicitly brings the non-extremal and extremal black holes at the same footing.

PACS numbers: 04070B, 0420

Isolated horizons [1], much like Killing horizons [2], were introduced in order to deal with the more practical problems involving black holes. Both formulations are *local*, in contrast with the global nature of event horizons [3]. However, isolated horizons, unlike Killing horizons, do not require a Killing vector field in their neighbourhood. Thus, isolated horizons are characterized by a weaker set of boundary conditions that give rise to both the zeroth and the first laws of black hole mechanics. Nevertheless, the isolated horizon boundary conditions, as developed in a series of papers [1, 4, 5, 6, 7, 8], consider a restricted equivalence class of null normal vector fields $[c\ell^a]$, where ℓ_1^a, ℓ_2^a belong to the class if and only if $\ell_1^a = c\ell_2^a$ for some positive definite constant c . Even though it has been emphasized that the most natural equivalence class of null normals at the horizon is $[\xi\ell^a]$, where ξ is an arbitrary positive function on the horizon, one restricts oneself, somewhat artificially perhaps, to the constant equivalence class. The purpose of this letter is to generalize the framework such that now the equivalence class of null normals is $[\xi\ell]$, where ξ is a given class of functions to be specified below, and to show how from this generalized setup both the zeroth and the first laws of black hole mechanics can be derived. This clearly makes isolated horizons applicable to a wider varieties of problems, both from analytical and numerical viewpoints. Another unexpected gain is that this generalized class of null normals include both non-extremal and extremal global solutions explicitly, namely there exists a choice of ξ for which the surface gravity associated with the vector field $\xi\ell^a$ vanishes, even though it is nonzero for other null normals in the class. In contrast, Killing horizons make essential use of the bifurcation two spheres in proving the zeroth and the first law [10], thereby distinguish extremal and non-extremal solutions in an explicit way.

Let $\mathcal{M}$ be a four-manifold equipped with a metric g_{ab} of signature $(-, +, +, +)$. Our notations and conventions closely follow that of [6]. Δ is a null hypersurface in $\mathcal{M}$ of which ℓ^a is a future directed null normal. However, if ℓ^a is a null normal, so is $\xi\ell^a$, where ξ is any positive

function on Δ . Thus, Δ naturally admits an equivalence class of null normals $[\xi\ell]$. We denote by $q_{ab} \triangleq g_{ab}$ the degenerate intrinsic metric on Δ induced by g_{ab} (indices that are not explicitly intrinsic on Δ will be pulled back and $\triangleq$ means that the equality holds *only on* Δ). A tensor q^{ab} will be an *inverse* of q_{ab} if it satisfies $q^{ab}q_{ac}q_{bd} \triangleq q_{cd}$. The expansion $\theta_{(\ell)}$ of the null normal ℓ is then defined by $\theta_{(\ell)} = q^{ab}\nabla_a\ell_b$, where ∇_a is the covariant derivative compatible with g_{ab} .

Boundary conditions and zeroth law: Δ is a non-expanding horizon (NEH) in $(\mathcal{M}, g_{ab})$ if the following conditions are satisfied: 1) Topologically $\Delta \equiv \mathbb{S}^2 \times \mathbb{R}$, 2) $\theta_{(\xi\ell)} \triangleq 0$ for any null normal $\xi\ell$ in the class $[\xi\ell]$ (actually, this is just *one* condition, since $\theta_{(\xi\ell)} \triangleq \xi\theta_{(\ell)}$, it suffices that $\theta_{(\ell)} \triangleq 0$), 3) all equations of motion including matter hold on Δ , in particular the matter energy-momentum tensor T_{ab} is such that $-T^a_b\xi\ell^b$ is future directed and causal on Δ . The important thing to note here is that all boundary conditions are intrinsic to Δ .

We shall always work with the null tetrad basis $(\ell, n, m, \bar{m})$ such that $1 = -n \cdot \ell = m \cdot \bar{m}$ and all other scalar products vanish. Since any null normal $\xi\ell$ is expansion-free, twist-free and geodesic, *i.e.* $\xi\ell^a\nabla_a(\xi\ell^b) \triangleq \kappa_{(\xi\ell)}\xi\ell^b$, where $\kappa_{(\xi\ell)}$ is the acceleration of $\xi\ell^a$, the Raychaudhuri equation becomes

$$0 \triangleq \mathcal{L}_{\xi\ell}\theta_{(\xi\ell)} \triangleq -|\sigma_{(\xi\ell)}|^2 - \xi^2 R_{ab}\ell^a\ell^b \quad (1)$$

where, $\sigma_{(\xi\ell)} = m^a m^b \nabla_a(\xi\ell_b)$ is the shear of $\xi\ell^a$. Since by energy condition $R_{ab}\ell^a\ell^b$ is positive both terms on the right hand side vanish independently on Δ . Therefore, every null normal $\xi\ell^a$ in the equivalence class is also shear-free.

From these results it follows that a class of one-forms $\omega_a^{(\xi\ell)}$ exists on Δ such that

$$\nabla_a(\xi\ell^b) \triangleq \omega_a^{(\xi\ell)}\xi\ell^b \quad (2)$$

The one-form varies over the class $[\xi\ell]$ as $\omega_a^{(\xi\ell)} = \omega_a^{(\ell)} + \nabla_a \ln \xi$. It also follows that every null normal $\xi\ell^a$ in the equivalence class is Killing on Δ , namely $\mathcal{L}_{(\xi\ell)}q_{ab} \triangleq 0$. It is useful to calculate the curvature of $\omega^{(\xi\ell)}$ as well. For the entire class of null-normals we find

$$d\omega^{(\xi\ell)} \triangleq 2(\text{Im}\Psi_2)\epsilon, \quad (3)$$

*Electronic address: ayan.chatterjee@saha.ac.in

†Electronic address: amit.ghosh@saha.ac.in

where $\text{Im}\Psi_2 = C_{abcd}\ell^a m^b \bar{m}^c n^d$ is a complex scalar constructed from the Weyl-tensor C_{abcd} and $\epsilon = im \wedge \bar{m}$.

The acceleration $\kappa_{(\xi\ell)}$ varies over the equivalence class $[\xi\ell]$ as follows: $\kappa_{(\xi\ell)} = \xi\kappa_{(\ell)} + \mathcal{L}_\ell \xi$, hence is not a constants on Δ . In order to obtain the zeroth law we need to restrict the NEH structure further. The restriction is called the *weak isolated horizon* (WIH) which is a NEH equipped with an equivalence class of null normals $[\xi\ell]$ obeying

$$\mathcal{L}_{(\xi\ell)} \omega^{(\xi\ell)} \triangleq 0. \quad (4)$$

Two comments are in order at this point. First of all, as in the case of Killing horizons, we will interpret the acceleration $\kappa_{(\xi\ell)}$ as the surface gravity associated with $\xi\ell^a$. However, a global Killing field being absent, the value of the surface gravity cannot be uniquely fixed. In isolated horizon formulation it is natural to maintain this freedom. Secondly, the boundary condition (4) defining a WIH is not a *single* condition representing the entire equivalence class, since $\mathcal{L}_{(\xi\ell)} \omega^{(\xi\ell)} \triangleq d(\xi\ell \cdot \omega^{(\xi\ell)}) \triangleq d\kappa_{(\xi\ell)}$ and $\kappa_{(\xi\ell)}$ is not constant for the entire class. To eliminate this pathology of having an infinite number of boundary conditions let us restrict our choice of the ξ -functions as follows: given say, $\kappa_{(\ell)} \triangleq \text{constant}$ on Δ , all $\kappa_{(\xi\ell)}$ are also constants on Δ if ξ satisfies the differential equation

$$\kappa_{(\xi\ell)} \triangleq \text{constant} \triangleq \xi\kappa_{(\ell)} + \mathcal{L}_\ell \xi. \quad (5)$$

Choosing a function v such that $\mathcal{L}_\ell v = 1$ one can solve the above equation for ξ obtaining $\xi = \eta e^{-v\kappa_{(\ell)}} + \kappa_{(\xi\ell)}/\kappa_{(\ell)}$, where η is a nonzero function satisfying $\mathcal{L}_\ell \eta = 0$. In the rest of the letter, we shall restrict ourselves to $\eta \triangleq \text{constant}$. With this restriction the boundary condition (4) becomes the *single* representative of the entire equivalence class. This class admits non-extremal as well as extremal global solutions, since ξ is positive even when $\kappa_{(\xi\ell)} \triangleq 0$.

As already noted, a direct consequence of the WIH boundary condition (4) is the *zeroth law*: $d\kappa_{(\xi\ell)} \triangleq 0$. Therefore, the surface gravity corresponding to each $\xi\ell$ in $[\xi\ell]$ is constant on Δ , provided ξ is restricted as above.

First Law: We consider the first order Palatini Lagrangian on $\mathcal{M}$ which has an internal boundary Δ . It is sufficient to consider $\mathcal{M}$ to be bounded by two partial Cauchy surfaces $M_\pm$, both oriented and cuts Δ at the two-spheres $S_\pm$ (see Figure 1). The basic fields are the co-tetrad e_a^I and the connection one-form A_{aI}^J . The lower case Latin letters are spacetime indices while the upper case letters refer to the internal four dimensional Minkowski spacetime having a fixed metric η_{IJ} of signature $(-+++)$. The metric on $\mathcal{M}$ is given by $g_{ab} = e_a^I e_b^J \eta_{IJ}$. Next we define a covariant derivative operator that acts only on internal indices

$$D_a v_I = \partial_a v_I + A_{aI}^J v_J, \quad (6)$$




FIG. 1: $M_\pm$ are two partial Cauchy surfaces enclosing a region of space-time and intersecting Δ in the 2-spheres $S_\pm$ respectively and extend to spatial infinity i^0 .

where, ∂ is a fiducial derivative operator chosen to be torsion free and compatible with the flat metric. We choose the Lagrangian four-form as

$$L = \frac{1}{16\pi G} [-\Sigma^{IJ} \wedge F_{IJ} + d(\Sigma^{IJ} \wedge A_{IJ})], \quad (7)$$

where $\Sigma^{IJ} = \frac{1}{2} \epsilon^{IJ}{}_{KL} e^K \wedge e^L$ and $F = dA + A \wedge A$.

Next consider the variation of (7). The allowed variations of the smooth fields (e_a^I, A) on $\mathcal{M}$ is such that it satisfies the standard conditions at infinity to ensure asymptotic flatness and on Δ must satisfy the following conditions: 1) each spacetime admits a null normal belonging to the equivalence class $[\xi\ell]$, 2) $(\Delta, [\xi\ell])$ is a WIH.

The standard variation gives on-shell $\delta L = d\Theta(\delta)$ where $16\pi G\Theta(\delta) = \delta\Sigma^{IJ} \wedge A_{IJ}$. Following standard procedures [9, 10], one then constructs the symplectic structure Ω on the space of solutions. First, we construct the symplectic current $J(\delta_1, \delta_2) = \delta_1\Theta(\delta_2) - \delta_2\Theta(\delta_1)$, which is closed on-shell. Integrating dJ over the spacetime under consideration we find

$$\int_{M_+ \cup M_- \cup \Delta \cup i^0} J = 0. \quad (8)$$

In order to evaluate $J|_\Delta$ we need expressions for the fields on Δ . Given the null vectors $(\ell, n, m, \bar{m})$ the tetrad e_a^I gives the internal null vectors $(\ell^I, n^I, m^I, \bar{m}^I)$. In terms of these vectors and with the help of the fact that the pull-back of the one-form $\ell_a \triangleq 0$, the pull-back of the two-form Σ^{IJ} can be expressed as

$$\begin{aligned} \Sigma_{\underline{ab}}^{IJ} \triangleq & 2\ell^{[I} n^{J]} \epsilon_{ab} + 2i(n_{[\underline{a}} m_{\underline{b}]} \ell^{[I} \bar{m}^{J]} \\ & - n_{[\underline{a}} \bar{m}_{\underline{b}]} \ell^{[I} m^{J]}). \end{aligned} \quad (9)$$

Using $\partial_a \ell_I = 0$ and the fact the ∇ is compatible with the tetrad e_a^I we can restrict the connection one-form A_{IJ} on Δ

$$A_{\underline{a}IJ} \triangleq -2\ell_{[I} n_{J]} \omega_a^{(\ell)} + \alpha_{\underline{a}IJ}, \quad (10)$$

where the one-form α_{IJ} satisfies the condition $\alpha_{IJ}\ell^J = 0$. This one-form decouples from rest of the analysis.

Formula (10) holds for the entire equivalence class $[\xi\ell]$ in the following precise sense: We have picked a fiducial null-normal ℓ from the class $[\xi\ell]$ as our basis such that the internal null vector ℓ_I is annihilated by the derivative ∂ and thus arrived at (10). But we could have chosen any other $\xi\ell$ as forming the basis $(\xi\ell, n/\xi, m, \bar{m})$. Carrying out the same analysis as before we would obtain the same decomposition (10).

In order to see that $J|_\Delta$ is an exact form we define a potential field $\psi_{(\xi\ell)}$ of the surface gravity $\kappa_{(\xi\ell)}$ as: $\mathcal{L}_{(\xi\ell)}\psi_{(\xi\ell)} \triangleq \kappa_{(\xi\ell)}$. Clearly $\psi_{(\xi\ell)}$ suffers from an additive ambiguity which can be fixed by requiring that $\psi_{(\xi\ell)}|_{S_-} \triangleq 0$. Using these definitions and the expressions (9) and (10) we find

$$\begin{aligned} J(\delta_1, \delta_2)|_\Delta &\triangleq \frac{1}{8\pi G} d[\delta_1\psi_{(\ell)}\delta_2\epsilon - \delta_2\psi_{(\ell)}\delta_1\epsilon] \\ &\triangleq: dj(\delta_1, \delta_2). \end{aligned} \quad (11)$$

Taking the orientation of the basis into account, the difference of the integrals $\int_M J - \oint_{S_\Delta} j$ is independent of the choice of the Cauchy surfaces M (S_Δ is the two-sphere at which M cuts Δ). Hence, one obtains the symplectic structure (see Figure 1)

$$\begin{aligned} \Omega(\delta_1, \delta_2) &= \frac{1}{16\pi G} \int_M [\delta_2\Sigma^{IJ} \wedge \delta_1 A_{IJ} - \delta_1\Sigma^{IJ} \wedge \\ &\delta_2 A_{IJ}] - \frac{1}{8\pi G} \oint_{S_\Delta} [\delta_2\epsilon\delta_1\psi_{(\ell)} - \delta_1\epsilon\delta_2\psi_{(\ell)}]. \end{aligned} \quad (12)$$

Given the symplectic structure the question now is whether the flow generated by the vector field $t^a = \xi\ell^a$ is Hamiltonian? To investigate this, we should calculate the one-form X_t where $X_t(\delta) = \Omega(\delta, \delta_t)$ and δ_t is to be interpreted as the Lie-flow $\mathcal{L}_t$ when acting on geometric fields. Then the necessary and sufficient condition for the existence of a Hamiltonian H_t is that the one-form X_t be exact $X_t = \mathbf{d}H_t$, where $\mathbf{d}$ is the exterior differential in the space of solutions. This makes $X_t(\delta) = \delta H_t$.

For the symplectic structure at hand, $X_t(\delta)$ receives contributions both from the bulk and the surface symplectic structures. The bulk term, thanks to the equations of motions satisfied by fields (e_I, A) and the linearized equations of motions satisfied by the variations $(\delta e_I, \delta A)$, contributes to the two-sphere boundaries of M , which are S_Δ and S_∞ . For the bulk term we get

$$X_t(\delta)|_M = -\frac{1}{8\pi G} \xi\kappa_{(\ell)}\delta\mathcal{A}_\Delta + \delta E_t \quad (13)$$

where, $\mathcal{A}_\Delta = \int_{S_\Delta} \epsilon$ is the area of S_Δ and E_t is the ADM-energy arising from the integral at S_∞ where it is understood that the asymptotic time translation is generated by t^a .

Care should be taken while calculating $X_t(\delta)$ from the surface symplectic structure, since one cannot identify $\delta_t\psi_{(\xi\ell)} = \mathcal{L}_t\psi_{(\xi\ell)}$. The reason is $\psi_{(\xi\ell)}$ being the potential of $\kappa_{(\xi\ell)}$, its variations are completely determined by

those of $\kappa_{(\xi\ell)}$. Since $\delta_t\kappa_{(\xi\ell)} = \mathcal{L}_t\kappa_{(\xi\ell)} \triangleq 0$ by zeroth law and $\delta_t\psi_{(\xi\ell)}|_{S_-} \triangleq 0$ it turns out that $\delta_t\psi_{(\xi\ell)} \triangleq 0$ everywhere. But $\psi_{(\xi\ell)} = \psi_{(\ell)} + \ln\xi$. Thus, $\delta_t\psi_{(\ell)} + \delta_t\ln\xi \triangleq 0$. From $\delta_t(\xi\ell^a) = \mathcal{L}_t(\xi\ell^a) = 0$ we find that $\delta_t\psi_{(\ell)} \triangleq -\mathcal{L}_t\xi$. Now we are in a position to calculate $X_t(\delta)$ arising from the surface symplectic structure

$$X_t(\delta)|_{S_\Delta} = -\frac{1}{8\pi G} \mathcal{L}_t\xi \delta\mathcal{A}_\Delta \quad (14)$$

Combining (13) and 14 we get

$$X_t(\delta) \triangleq -\frac{1}{8\pi G} \kappa_{(\xi\ell)}\delta\mathcal{A}_\Delta + \delta E_t. \quad (15)$$

This gives the first-law $\delta E_\Delta^t = \kappa_{(\xi\ell)}\delta\mathcal{A}_\Delta/8\pi G$ once we identify $X_t = \mathbf{d}H_t$ and define $E_t - H_t = E_\Delta^t$, where E_Δ^t is the horizon-mass. That X_t is a closed form forces one to regard $\kappa_{(\xi\ell)}$ as a function of the area $\mathcal{A}_\Delta$ alone.

Electromagnetic Field: NEH boundary conditions restricts the matter energy-momentum tensor by $T_{ab}\ell^a\ell^b \triangleq 0$. For the electromagnetic fields $\mathbf{A}$ this implies

$$0 \triangleq T_{ab}\ell^a\ell^b \triangleq |\ell^a m^b \mathbf{F}_{ab}|^2, \quad (16)$$

where $\mathbf{F} = d\mathbf{A}$ is the electromagnetic field strength. It follows from (16) that $\ell^a \mathbf{F}_{a\bar{b}} \triangleq 0$ and $\ell^a * \mathbf{F}_{a\bar{b}} \triangleq 0$. These two restrictions tell us there is no flux of electromagnetic radiation *across* the horizon. Radiation may flow along the horizon. Since Δ is an inner boundary and the normal to S_Δ is inward pointing, the electric charge of the horizon is defined by (it is assumed that all magnetic charges are zero)

$$Q_\Delta \triangleq -\frac{1}{4\pi} \oint_{S_\Delta} * \mathbf{F}. \quad (17)$$

Since $\mathcal{L}_{(\xi\ell)} * \mathbf{F} \triangleq 0$ the charge Q_Δ is independent of the choice of S_Δ .

Zeroth Law: To establish the zeroth law for the electromagnetic case, we need to define an electric potential Φ at the horizon. For this the electromagnetic potential $\mathbf{A}$ is gauge fixed at the WIH such that

$$\mathcal{L}_{(\xi\ell)} \mathbf{A}_{\underline{a}} \triangleq \nabla_{\underline{a}} \mathcal{L}_{(\xi\ell)} \chi_{(\xi\ell)}. \quad (18)$$

where $\chi_{(\xi\ell)}$ is a fixed function of v . Given such an electromagnetic potential $\mathbf{A}$ we can now define the scalar potential $\Phi_{(\xi\ell)}$ at the horizon as

$$\Phi_{(\xi\ell)} \triangleq -\xi\ell \cdot \mathbf{A} + \mathcal{L}_{(\xi\ell)} \chi_{(\xi\ell)}. \quad (19)$$

From (18) it follows immediately that $d\Phi_{(\xi\ell)} \triangleq 0$, hence $\Phi_{(\xi\ell)}$ is constant on the horizon. In order that the potential be constant for the entire equivalence class $[\xi\ell]$ requires the gauge fixing functions to be related as $\mathcal{L}_{(\xi\ell)}\chi_{(\xi\ell)} - \Phi_{(\xi\ell)} = \xi[\mathcal{L}_\ell\chi_{(\ell)} - \Phi_{(\ell)}]$.

First Law: The Lagrangian four-form for the electromagnetic field is given by $8\pi L = -\mathbf{F} \wedge \star \mathbf{F}$. The variation for the Lagrangian is carried out such that $\mathbf{A}$ has the expected asymptotic fall-off and is gauge fixed at Δ as in (18). Here the key point is that although we needed a gravitational surface term in the action, thanks to the electromagnetic zeroth law, such a term is not needed here. Proceeding as in the gravitational case here also we find a bulk and a surface term in the symplectic structure. The surface term can be made explicit by introducing a potential for $\Phi_{(\xi\ell)}$: $\mathcal{L}_{(\xi\ell)}\varphi_{(\xi\ell)} \triangleq -\Phi_{(\xi\ell)}$. It too suffers from an additive ambiguity which is removed by choosing $\varphi_{(\xi\ell)}|_{S_-} \triangleq 0$. Then the symplectic structure is

$$\Omega_{\text{em}}(\delta_1, \delta_2) = -\frac{1}{4\pi} \int_M [\delta_1 \star \mathbf{F} \wedge \delta_2 \mathbf{A} - (1 \leftrightarrow 2)] + \frac{1}{4\pi} \oint_{S_\Delta} [\delta_1 \star \mathbf{F} \delta_2 (\chi_{(\xi\ell)} + \varphi_{(\xi\ell)}) - (1 \leftrightarrow 2)] . \quad (20)$$

Again, we look for evaluating $X_t(\delta)$ from the electromagnetic part of the symplectic structure. Making use of the field equations the bulk symplectic structure receives contributions from the boundary alone, which is found to be $(\mathcal{L}_t \chi_{(\xi\ell)} - \Phi_{(\xi\ell)}) \delta Q_\Delta$. To evaluate the contribution from the surface symplectic structure care must be taken not to identify δ_t with $\mathcal{L}_t$ for the derived potential φ . It

turns out that $\delta_t \varphi_{(\xi\ell)} \triangleq 0$ everywhere and the nonzero contribution is $-\delta_t \chi_{(\xi\ell)}$. Combining the bulk and the surface contributions we find that

$$X_t^{\text{em}}(\delta) \triangleq -\Phi_{(\xi\ell)} \delta Q_\Delta . \quad (21)$$

In this letter we sketched how the boundary conditions defining a WIH can be relaxed for a larger equivalence class of null normals so that still the zeroth and the first laws of black hole mechanics emerge. The framework should be extendible to include rotations as well. Since this formulation is blind to extremal and non-extremal global solutions it is believed that the entropy issue can be addressed more squarely if one quantizes this WIH. Generalizations to dynamical horizons and their settling down to such WIH would also be addressed in the future.

Acknowledgments

We thank Abhay Ashtekar for discussion and clarification of a point from the reference [6]. We thank Ashok Chatterjee for discussion. We especially thank Parthasarathi Majumdar for several stimulating discussions and continuous encouragement during the course of the work.

-
- [1] A. Ashtekar, C. Beetle, O. Dreyer, S. Fairhurst, B. Krishnan, J. Lewandowski and J. Wisniewski, *Phys. Rev. Lett.* **85**, 3564 (2000).
 - [2] B. S. Kay and R. M. Wald, *Phys. Rept.* **207**, 49 (1991).
 - [3] J.W. Bardeen, B. Carter and S.W. Hawking, *Commun. Math. Phys.* **31** 161 (1973).
 - [4] A. Ashtekar, C. Beetle and S. Fairhurst, *Class. Quantum Grav.* **16** L1 (1999).
 - [5] A. Ashtekar, C. Beetle and S. Fairhurst, *Class. Quantum Grav.* **17** 253 (2000).
 - [6] A. Ashtekar, S. Fairhurst and B. Krishnan, *Phys. Rev. D* **62** 104025 (2000).
 - [7] A. Ashtekar, C. Beetle and J. Lewandowski, *Phys. Rev. D* **64** 044016 (2001).
 - [8] A. Ashtekar, C. Beetle and J. Lewandowski, *Class. Quantum Grav.* **19** 1195 (2002).
 - [9] A. Ashtekar, L. Bombelli and O. Reula, *Mechanics, Analysis and Geometry: 200 years after Lagrange* ed. M Francaviglia (North-Holland) (1991).
 - [10] R. M. Wald, *Phys. Rev. D* **48** 3427 (1993).